arXiv:1207.2822v1 [math.AT] 12 Jul 2012

ON KIRCHHOFF'S THEOREMS WITH COEFFICIENTS IN A LINE BUNDLE

MICHAEL J. CATANZARO, VLADIMIR Y. CHERNYAK, AND JOHN R. KLEIN

Abstract. We prove 'twisted' versions of Kirchhoff's network theorem and Kirchhoff's matrix-tree theorem on connected finite graphs. Twisting here refers to chains with coefficients in a flat unitary line bundle.

Contents

1. Introduction 1
2. Properties of twisted spanning trees 7
3. Proof of Theorem A and Corollary B 11
4. Proof of Theorem C 13
References 15

1. Introduction

It is well-known that the classical result of Kirchhoff on the flow of electricity through a finite network admits an elegant formulation using algebraic topology [E],[NS],[R]. For a finite connected 1-dimensional CW complex Γ, the real cellular chain complex $\partial\colon C_1(\Gamma;\mathbb{R}) \to C_0(\Gamma;\mathbb{R})$ is a homomorphism of finite dimensional real inner product spaces with orthonormal basis given by the set of cells. When the branches of the network have unit resistance, Kirchhoff's *network theorem* is reflected in the statement that the restricted homomorphism

$$\partial\colon B^1(\Gamma;\mathbb{R}) \to B_0(\Gamma;\mathbb{R}) \tag{1}$$

is an isomorphism, where $B_0(\Gamma;\mathbb{R})$ is the vector subspace of zero-boundaries and $B^1(\Gamma;\mathbb{R})$ is the orthogonal complement to the space of 1-cycles $Z_1(\Gamma;\mathbb{R}) \subset C_1(\Gamma;\mathbb{R})$ (when the network has branches of varying resistance, one rescales the inner product on $C_1(\Gamma;\mathbb{R})$ accordingly).

Date: July 13, 2012.
2010 *Mathematics Subject Classification.* Primary: 55J15, 57M15, 57Q10 Secondary: 05C05, 05C21, 05E45, 82C31.

Actually Kirchhoff's network theorem does more in that it provides a concrete expression for the inverse to the isomorphism (1) in terms of the set of spanning trees of Γ. The expression amounts to an explicit formula for the orthonormal projection of $C_1(\Gamma;\mathbb{R})$ onto $Z_1(\Gamma;\mathbb{R})$ in terms of the set of spanning trees of Γ.

A companion result of Kirchhoff, which has gotten more press, is the *matrix-tree theorem*, which computes the determinant of the restricted combinatorial Laplacian

$$\partial\partial^* \colon B_0(\Gamma;\mathbb{R}) \to B_0(\Gamma;\mathbb{R})\,.$$

Here, ∂^* denotes the formal adjoint to the boundary operator (for a slightly different formulation, see [B, p. 57]). In the unit resistance case, the result says that that $\det(\partial\partial^*)$ equals the product of the number of vertices with the number of spanning trees of Γ. In [CCK], using ideas from statistical mechanics, we showed how the matrix-tree theorem can be derived from the general case of the network theorem. We also generalized both of these results to higher dimensional CW complexes.

The main purpose of the current paper is to derive a twisted version of Kirchhoff's theorems. Here the twisting is given by taking coefficients in a *complex line bundle*. In physical terms, the twisted version of the network theorem turns out to model the flow of current through an electrical network in the presence of *fluctuations*.[1] As an application, we deduce a twisted version of the matrix-tree theorem. Suitably reformulated, our twisted matrix-tree theorem is actually a result of Forman [F, eq. (1)] which we first learned about in a recent paper of Kenyon [Ke, thm. 5]. Forman's proof is combinatorial, using an explicit expression for the determinant in terms of symmetric groups. Kenyon's proof relies on the Cauchy-Binet theorem. By contrast, our approach is inspired by statistical mechanical ideas and closely follows the untwisted version appearing in [CCK].

Graphs. A *graph* Γ is a CW complex of dimension one. We let Γ_0 denote the set of 0-cells and Γ_1 the set of 1-cells. A 0-cell is called a *vertex* and a 1-cell is called an *edge*. The entire structure of Γ is given by a function

$$(d_0, d_1)\colon \Gamma_1 \to \Gamma_0 \times \Gamma_0\,, \tag{2}$$

which sends an edge b to its initial and terminal endpoints (where the edge b is oriented using its characteristic map $\chi_b\colon [0,1] \to \Gamma$). Given the function (2), one can reconstruct Γ by taking

$$\Gamma_0 \cup (\Gamma_1 \times [0,1])$$

[1] We will develop the physical interpretation in another paper.

where the union is amalgamated over the map $\Gamma_1 \times \{0,1\} \to \Gamma_0$ given by $(b,0) \mapsto d_0(b)$ and $(b,1) \mapsto d_1(b)$.

A *loop edge* is an edge such that $d_0(b) = d_1(b)$. If b is not a loop edge, it is said to be *regular*.

Flat line bundles on graphs. A *flat complex vector bundle* ρ on a graph Γ is a rule which assigns to each vertex $i \in \Gamma_0$ a finite rank complex vector space V_i over $\mathbb{C}$ and to each edge b with $(d_0 b, d_1 b) = (i,j)$ an isomorphism

$$\rho_b \colon V_i \to V_j \,.$$

We say that ρ is *unitary* if each V_i is a hermitian inner product space and each ρ_b is unitary. In this paper we will deal exclusively with the rank one case, i.e., flat complex line bundles. Henceforth, we simplify terminology and refer to ρ as a *line bundle*.[2]

Given a line bundle ρ, by choosing a non-zero vector u_i in V_i having unit norm, we can identify V_i with $\mathbb{C}u_i$, the complex vector space spanned by u_i. Consequently, there is no loss in generality in assuming that $V_i = \mathbb{C}$ for every $i \in \Gamma_0$. In this instance $\rho_b \colon \mathbb{C} \to \mathbb{C}$ is given by multiplication by a unit complex number, which by abuse of notation we denote as ρ_b. With respect to these choices, ρ is given by a function $\Gamma_1 \to U(1)$.

Recall that a circuit C of Γ is a simple closed path. An orientation of C consists of a choice of direction for traversing C.

Definition 1.1. If C is an oriented circuit of Γ, then the *holonomy* of ρ along C is given by the product

$$\rho_C := \prod_{b \in C_1} \rho_b^{s_b}$$

where $s_b = \pm 1$ according as to whether the orientation of C is the same as the orientation of b.

If $\bar{C}$ denotes C with its reverse orientation, then $\rho_{\bar{C}} = \rho_C^*$, where ρ_C^* denotes the complex conjugate of ρ_C.

More generally, if $A \subset \Gamma$ is a subgraph, in which each circuit C of A is given an orientation, then we write

$$\rho_A = \prod_C \rho_C$$

where the product ranges over the circuits C of A.

[2]Strictly speaking, what we have defined here is really the notion of a transport operator on Γ associated with a flat connection, but we will not need to worry about this distinction.

Lastly, we set

$$\hat{\rho}_A = \prod_C (\rho_C - 1)(\rho_C^* - 1) = \prod_C (2 - \rho_C - \rho_C^*)$$

where C ranges over the circuits of A. This last expression is independent of the choice of orientation for the circuits and is also a real number.

The twisted chain complex. For $i = 0, 1$, let $C_i(\Gamma; \rho)$ denote the $\mathbb{C}$-vector space having basis Γ_i. Define the twisted boundary operator

$$\partial \colon C_1(\Gamma; \rho) \to C_0(\Gamma; \rho)$$

by mapping an edge b to the vector $\rho_b d_0(b) - d_1(b)$ and extending linearly. The homology of this two-stage complex is denoted by $H_*(\Gamma; \rho)$. It is invariant with respect to barycentric subdivision. That is, if Γ' is the barycentric subdivision of Γ and ρ' is a line bundle on Γ' such that $\rho_b = \rho'_{b_0}\rho'_{b_1}$ when $b = b_0 b_1$ is the subdivision of an edge b, then $H_*(\Gamma; \rho) \cong H_*(\Gamma'; \rho')$. Note that $H_1(\Gamma; \rho)$ is a subspace of $C_1(\Gamma; \rho)$ consisting of the cycles.

Remark 1.2. This is the traditional notation. It is imprecise since each vector space $C_i(\Gamma; \rho)$ does not depend on ρ whereas the boundary operator ∂ does. A more precise notation would write the complex as $\partial_\rho \colon C_1(\Gamma; \mathbb{C}) \to C_0(\Gamma; \mathbb{C})$.

If $A \subset \Gamma$ is a subcomplex, we have the relative chain complex $C_*(\Gamma, A; \rho)$ which is the quotient complex $C_*(\Gamma; \rho)/C_*(A; \rho)$. It has a basis consisting of the cells of Γ which are not in A.

The resistance operator. A *resistance function* is a map $r \colon \Gamma_1 \to \mathbb{R}_+$ which assigns to an edge b a resistance $r_b > 0$. Associated with r is the *resistance operator*

$$R \colon C_1(\Gamma; \rho) \to C_1(\Gamma; \rho)\,,$$

which on basis elements is defined by $b \mapsto r_b b$.

The standard and modified inner products. The *standard Hermitian inner product* on $C_1(\Gamma; \rho)$, denoted $\langle\ ,\ \rangle$, is given on basis elements $b, b' \in \Gamma_1$ by

$$\langle b, b' \rangle := \delta_{bb'}\,,$$

where $\delta_{bb'}$ is Kronecker delta.

Associated with the resistance operator R is the *modified inner product* on $C_1(\Gamma; \rho)$, denoted $\langle\ ,\ \rangle_R$, is given by

$$\langle b, b' \rangle_R := r_b \langle b, b' \rangle = \delta_{bb'} r_b\,.$$

Twisted spanning trees. From now on we assume that Γ is connected and finite.

Definition 1.3. A *ρ-spanning tree* for Γ is a subcomplex $T \subset \Gamma$ such that

- $T_0 = \Gamma_0$,
- $H_1(T;\rho) = 0$, and
- $H_0(T;\rho) \to H_0(\Gamma;\rho)$ is an isomorphism.

Remark 1.4. When ρ is the trivial line bundle, we recover the usual notion of spanning tree. In the next section we characterize the ρ-spanning trees of Γ.

Definition 1.5. The *weight* of a ρ-spanning tree T is the real number

$$w_T := \hat{\rho}_T \prod_{b \in T_1} r_b^{-1} .$$

The operator $\bar{T}$. Given a ρ-spanning tree T, we define an operator

$$\bar{T} \colon C_1(\Gamma;\rho) \to H_1(\Gamma;\rho)$$

as follows: if $b \in T_1$ then $\bar{T}(b) = 0$. If $b \in \Gamma_1 \setminus T_1$, we form the graph $T \cup b$. Then $\dim_{\mathbb{C}} H_1(T \cup b;\rho) = 1$. Let $c \in H_1(T \cup b;\rho)$ be a non-zero vector, and set $t_b = \langle c, b\rangle$. We set $\bar{T}(b) := c/t_b$. This does not depend on the choice of c. Note that $H_1(T \cup b;\rho) \to H_1(\Gamma;\rho)$ is an inclusion, so this definition makes sense.

Remark 1.6. It will be useful to have an alternative description of $\bar{T}$. Assume $b \in \Gamma_1 \setminus T_1$. The homology class $[\partial b] \in H_0(T;\rho) = 0$ is trivial, so $\partial b \in C_0(T;\rho)$ bounds a chain $u \in C_1(T;\rho)$. Then $c := b - u \in C_1(T \cup b;\rho)$ is a cycle such that $t_b = \langle c, b\rangle = 1$. In this case $\bar{T}(b) = c$.

The main results. The twisted version of Kirchhoff's network theorem will be a consequence of having a concrete description of the projection operator from twisted 1-chains to twisted 1-cycles.

Theorem A (Twisted Projection Formula)**.** *With respect to the modified inner product $\langle\ ,\ \rangle_R$, the hermitian projection of $C_1(\Gamma;\rho)$ onto the subspace $H_1(\Gamma;\rho)$ is given by*

$$\tfrac{1}{\Delta} \sum_T w_T \bar{T} ,$$

where T ranges over the ρ-spanning trees of Γ and $\Delta = \sum_T w_T$.

Our twisted version of Kirchhoff's network theorem is

Corollary B (Twisted Network Theorem)**.** *Given a vector* $\mathbf{V} \in C_1(\Gamma;\rho)$*, there is only one vector* $z \in Z_d(\Gamma;\rho)$ *such that* $\mathbf{V} - Rz \in B^1(\Gamma;\rho)$*. Furthermore, for each edge* b*, we have*

$$\langle z, b\rangle = \tfrac{1}{\Delta}\sum_T \tfrac{w_T}{r_b}\langle \mathbf{V}, \bar{T}(b)\rangle\,.$$

Remark 1.7. In the untwisted case ($\rho = 1$), this is the formulation of Kirchhoff's network theorem that is found in [NS]. The expression $\langle \mathbf{V}, b\rangle$ is called the voltage source on the branch b and $\langle z, b\rangle$ is the current residing on b.

Let

$$\partial_R^*\colon C_0(\Gamma;\rho) \to C_1(\Gamma;\rho)$$

be the formal adjoint to the boundary operator $\partial\colon C_1(\Gamma;\rho) \to C_0(\Gamma;\rho)$ with respect to the standard hermitian inner product on $C_0(\Gamma;\rho)$ and the modified one on $C_1(\Gamma;\rho)$ as determined by the resistance operator R.

The following is the result of Forman [F, eq. (1)] that was alluded to above.

Theorem C (Twisted Weighted Matrix-Tree Theorem)**.**

$$\det(\partial\partial_R^*\colon C_0(\Gamma;\rho) \to C_0(\Gamma;\rho)) = \sum_T w_T\,,$$

where T *ranges over all* ρ*-spanning trees, and* w_T *is as in Definition 1.5.*

Remark 1.8. When $H_0(\Gamma;\rho)$ non-trivial, the determinant $\det(\partial\partial_R^*)$ vanishes. We show in the next section that the triviality of $H_0(\Gamma;\rho)$ is equivalent to the existence of a ρ-spanning tree. Hence, the sum $\sum_T w_T$ is trivial when $H_0(\Gamma;\rho)$ is non-trivial, and Theorem C holds trivially in this case. Consequently, from now on we may assume that $H_0(\Gamma;\rho)$ is trivial.

The case $R = 1$ is worth singling out. We use the notation $\partial^* = \partial_R$ in this case.

Corollary D (Twisted Matrix-Tree Theorem)**.**

$$\det(\partial\partial^*) = \sum_T \hat{\rho}_T\,,$$

where T *ranges over all* ρ*-spanning trees.*




FIG. 1. A lattice graph equipped with CRSF having four components (cf. Remark 2.2). The edges of the graph are the gray dashed lines. The edges of the CRSF are indicated in black.

Outline. In § 2 we develop foundational material on ρ-spanning trees. § 3 contains the proofs of Theorem A and Corollary B. In § 4, we prove Theorem C using Theorem A and the low temperature limit argument of [CCK].

Acknowledgements. The authors wish to Nikolai Sinitsyn for discussions related to the physical interpretation of the twisted network theorem. We are also indebted Misha Chertkov for hosting the first author and to Andrei Piryatinski for his unlimited hospitality. We thank the Los Alamos Center for Nonlinear Studies and the T-4 division for partially supporting this research. This material is based upon work supported by the National Science Foundation under Grant Nos. CHE-1111350 and DMS-1104355.

2. Properties of twisted spanning trees

The following lemma characterizes the structure of ρ-spanning trees of Γ.

Lemma 2.1. *Assume $H_0(\Gamma;\rho) = 0$. A subcomplex $T \subset \Gamma$ is a ρ-spanning tree if and only if*

- $T_0 = \Gamma_0$,
- *each connected component T^α of T has trivial Euler characteristic, i.e., T^α possesses a unique circuit, C_α and*
- *the holonomy around C_α is non-trivial.*

Remark 2.2. A *cycle-rooted spanning forest (CRSF)* is a subcomplex T of Γ satisfying the first two conditions listed in Lemma 2.1, i.e., a ρ-spanning tree is a CRSF additionally satisfying the condition that the holonomy around circuits is non-trivial (cf. Fig. 1, [Ke, 4.1]).

Proof of Lemma 2.1. Assume T is a ρ-spanning tree. The assumption $H_0(\Gamma;\rho) = 0$ implies that $H_0(T;\rho) = 0$ and the latter implies

$H_0(T^\alpha;\rho) = 0$ since $H_0(T;\rho) = \oplus_\alpha H_0(T_\alpha;\rho)$. Similarly, $H_1(T;\rho) = 0$ implies $H_1(T^\alpha;\rho) = 0$. Hence the chain complex

$$\partial\colon C_1(T^\alpha;\rho) \to C_0(T^\alpha;\rho)$$

is acyclic. In particular, the number of edges of T^α equals the number of vertices, so the Euler characteristic of T^α is trivial. Orient the unique circuit C_α and let the holonomy around C_α be denoted ρ_α. Then independence of twisted cohomology with respect to subdivision yields $H_0(T^\alpha;\rho) = H_0(C_\alpha;\rho) = H_0(S^1;\rho_\alpha)$, where we are thinking of S^1 as a graph with one vertex and one edge and where the line bundle is given by ρ_α. An easy calculation shows $H_0(S^1;\rho_C)$ is the cokernel of the map $(\rho_\alpha - 1)\cdot\colon \mathbb{C} \to \mathbb{C}$. Hence the triviality of $H_*(T^\alpha;\rho_\alpha)$ is equivalent to the statement $\rho_\alpha \neq 1$.

Conversely, given T satisfying the three conditions, the second and third conditions imply $H_*(T;\rho) = \oplus_\alpha H_*(T^\alpha;\rho)$ is trivial. Hence T is a ρ-spanning tree. □

Lemma 2.3. *Assume $H_0(\Gamma;\rho) = 0$. Then Γ has a ρ-spanning tree.*

Proof. Call an edge b of Γ *essential* if there is a cycle $z \in H_1(\Gamma;\rho) \subset C_1(\Gamma;\rho)$ such that $\langle b, z\rangle \neq 0$. If there is no such edge, then it is straightforward to check that Γ is a ρ-spanning tree.

Assume then that there is an essential edge b. Let Y be the effect of removing (the interior of) b from Γ. Then we have a short exact sequence

$$0 \to H_1(Y;\rho) \to H_1(\Gamma;\rho) \to H_1(b,\partial b;\rho) \to H_0(Y;\rho) \to 0$$

and the condition $\langle b, z\rangle \neq 0$ implies that the homomorphism $H_1(\Gamma;\rho) \to H_1(b,\partial b;\rho)$ is non-trivial (note that $H_1(b,\partial b;\rho) \cong \mathbb{C}$). It follows that $\dim_{\mathbb{C}} H_1(Y;\rho) < \dim_{\mathbb{C}} H_1(\Gamma;\rho)$ and $H_0(Y;\rho) = 0$. We do not require that Y be connected. We now replace Γ by Y and iterate this construction until we obtain a subcomplex T having no essential cells and $H_*(T;\rho) = 0$. Then T is a ρ-spanning tree. □

Lemma 2.4. *Fix a ρ-spanning tree T and let $b_1, \ldots, b_k$ be the set of edges of $\Gamma_1 \setminus T_1$. Then $\{\bar{T}(b_1), \cdots, \bar{T}(b_k)\}$ is a basis for $H_1(\Gamma;\rho)$.*

Proof. The homomorphism $H_1(\Gamma;\rho) \to H_1(\Gamma, T;\rho)$ an isomorphism. Furthermore, $H_1(\Gamma, T;\rho) = C_1(\Gamma;T;\rho)$ has basis $\{b_1, \cdots, b_k\}$. The inverse homomorphism sends b_i to $\bar{T}(b_i)$. □

Corollary 2.5. *For any $z \in H_1(\Gamma;\rho)$, we have $\bar{T}(z) = z$.*

Proof. The definition of $\bar{T}$ shows $\bar{T}^2(b_i) = \bar{T}(b_i)$. Write $z = \sum_i a_i \bar{T}(b_i)$. Then

$$\bar{T}(z) = \sum_i a_i \bar{T}^2(b_i) = \sum_i a_i \bar{T}(b_i) = z\,.$$ □

Given a ρ-spanning tree T, consider an edge $b_i \in \Gamma_1 \setminus T_1$ as well as an edge $b_j \in T_1$. Let $U = (T \setminus b_j) \cup b_i$.

Lemma 2.6. *U is a ρ-spanning tree if and only if $\langle \bar{T}(b_i), b_j \rangle \neq 0$.*

Proof. Throughout this proof we use local coefficients in ρ but suppress this from the notation. We have an exact sequence

$$0 \to H_1(T \setminus b_j) \to H_1(T) \to H_1(b_j, \partial b_j) \to H_0(T \setminus b_j) \to 0\,,$$

where we are using the fact that $H_0(T) = 0$. Since $H_1(T) = 0$ and $\dim_{\mathbb{C}} H_1(b_j, \partial b_j) = 1$, we infer that $H_1(T \setminus b_j) = 0$ and $\dim_{\mathbb{C}} H_0(T \setminus b_j) = 1$.

The inclusion $U \subset T \cup b_i$ induces another exact sequence

$$0 \to H_1(U) \to H_1(T \cup b_i) \to H_1(b_j, \partial b_j) \to H_0(U) \to 0\,,$$

and the homomorphism $H_1(T \cup b_i) \to H_1(b_j, \partial b_j)$ is a map of rank one vector spaces that is induced by sending the preferred cycle $c \in H_1(T \cup b)$ to $\langle \bar{T}(b_i), b_j \rangle$ with respect to the preferred identification $H_1(b, \partial b) \cong \mathbb{C}$. Consequently, U is a ρ-spanning tree if and only if $\langle \bar{T}(b_i), b_j \rangle \neq 0$. □

Proposition 2.7. *With b_i, b_j, T, U as above, we have*

$$\hat{\rho}_T \langle \bar{T}(b_i), b_j \rangle = \hat{\rho}_U \langle b_i, \bar{U}(b_j) \rangle\,.$$

Proof. Case 1: Assume that b_i is attached to two distinct components of T, say A and B. By switching the roles of A and B if necessary, we may suppose that $(d_0(b_i), d_1(b_i)) = (w, v)$, where v lies in A and w lies in B. Without loss in generality assume that b_j lies in A. Let C be the unique circuit of A and C' the unique circuit of B.

Then

$$v = \partial c$$

for $c \in C_1(A; \rho)$. We may then write $c = c_0 + \alpha b_j$, where $\alpha \in \mathbb{C}$ and $\langle c_0, b_j \rangle = 0$. Similarly, we write $w = \partial d$, where $d \in C_1(B; \rho)$. Then $\langle \bar{T}(b_i), b_j \rangle$ equals

$$\langle b_i - (\rho_{b_i} d - c_0 - \alpha b_j), b_j \rangle = \alpha\,,$$

since $\partial(\rho_{b_i} d - c_0 - \alpha b_j) = \partial b_i$ and $\rho_{b_i} d - c_0 - \alpha b_j$ is a chain of T.

A similar calculation shows $\langle b_i, \bar{U}(b_j) \rangle$ equals

$$\langle b_i, b_j - \tfrac{\rho_{b_i} d - b_i - c_0}{\alpha} \rangle = (\alpha^{-1})^*\,.$$

In order to compute α, it is enough to identify the 1-chain $c \in C_1(A; \rho)$ whose boundary equals v, since then $\langle c, b_j \rangle = -\alpha$.

To find c we rename $v = v_1$ and choose a vertex v_k on the unique circuit of A together with an embedded path of edges $e_1, \ldots e_k$ which

connects v_1 to v_k. Without loss in generality, we can assume that none of the edges e_i lies in the unique cycle of A. Let $e_{k+1}, \dots, e_n$ denote the sequence of edges given by the traversing the unique cycle of A such that v_k is a vertex of both e_{k+1} and e_n. Then c is a linear combination of the edges e_i which can be explicitly computed using the fact that $\partial e_i = \rho_{e_i} d_0(e_i) - d_1(e_i)$. Then a straightforward calculation yields the expression for the component of c along the edge e_i as

$$\langle c, e_i \rangle = -\frac{\rho_1^{s_1} \cdots \rho_{i-1}^{s_{i-1}}}{\rho_A - 1}\,, \tag{3}$$

where $\rho_i := \rho_{e_i}$ and $s_i = \pm 1$ according as to whether e_i points in the direction of the path or not (we have also oriented A in a way that is compatible with our choice of path). In particular, $b_j = e_\ell$ for some index ℓ, so

$$\alpha = \frac{\rho_1^{s_1} \cdots \rho_{\ell-1}^{s_{\ell-1}}}{\rho_A - 1}\,,$$

Since $\hat{\rho}_T = \hat{\rho}_A \hat{\rho}_B \hat{\rho}'$, where $\hat{\rho}'$ is the product of the $\hat{\rho}_{T^\alpha}$ ranging over the remaining components of T, we have

$$\hat{\rho}_T \alpha = \rho_1^{s_1} \cdots \rho_{i-1}^{s_{\ell-1}} (\rho_A^* - 1) \hat{\rho}_B \hat{\rho}'\,.$$

Since $\hat{\rho}_U = \hat{\rho}_B \hat{\rho}'$ we see

$$\hat{\rho}_U (\alpha^{-1})^* = \hat{\rho}_T \alpha\,,$$

which concludes Case 1.

Case 2: In this instance b_i is attached to a single component A of T. In this case we need to find a 1-chain c of A such that $\partial c = \partial b_i$. Arguing in an analogous way as in the beginning of Case 1, if we set

$$\langle \bar{T}(b_i), b_j \rangle = \alpha \tag{4}$$

then it follows that

$$\langle \bar{b}_i, \bar{U}(b_j) \rangle = (\alpha^{-1})^*\,. \tag{5}$$

Suppose $d_1(b_i) = v$ and $d_0(b_i) = w$ (where it is possible that $v = w$). We select a simple path $e_1, \cdots, e_k$ of edges of A such that v meets e_1 and w meets e_k. Let us rename b_i as e_{k+1}. Then c is a linear combination of the edges e_i for $1 \leq i \leq k$. Let C denote the unique circuit of A. Then A decomposes as

$$A_- \cup C \cup A_+$$

in which A_- is a connected subgraph of A that meets the vertex v and A_+ is the connected subgraph of A which meets the vertex w. For a

given index i, consider the expressions

$$\alpha_- := \prod_{1\le j<i} \rho_j^{s_j} \quad \alpha_+ := \prod_{i\le j\le k+1} \rho_j^{s_j}\,.$$

A calculation similar to that appearing in Case 1 gives, for $0 \le i \le k$,

$$\langle c, e_i\rangle = \begin{cases} \alpha_-\,, & \text{if } e_i \subset A_-\,,\\ (\rho_A - 1)^{-1}(\alpha_+ - \alpha_-)\,, & \text{if } e_i \subset C\,,\\ \alpha_+\,, & \text{if } e_i \subset A_+\,. \end{cases} \tag{6}$$

Then if $\beta_j = e_\ell$ for some ℓ, we have $\langle \bar{T}(\beta_i), \beta_j\rangle = -\langle c, e_\ell\rangle$. As before, we have $\hat{\rho}_T = \hat{\rho}_A\hat{\rho}'$, where $\hat{\rho}'$ is a product of $\hat{\rho}_{T^\alpha}$ for T^α ranging over the other components of T. Consequently,

$$\langle \hat{\rho}_T\bar{T}(\beta_i), \beta_j\rangle = \hat{\rho}_A\hat{\rho}'\alpha$$

where $\alpha = -\langle c, e_\ell\rangle$ is explicitly given by Eqn. (6).

The remainder of the argument is just as in Case 1. A straightforward calculation that we omit shows

$$\hat{\rho}_U = \begin{cases} \hat{\rho}_T & b_j \subset A_- \cup A_+\,,\\ (2 + \alpha_+\alpha_-^* - \alpha_+^*\alpha_-)\hat{\rho}' & b_j \subset C \end{cases} \tag{7}$$

Then use Eq. (5) and Eq. (6) to identify the product $\hat{\rho}_U(\alpha^{-1})^*$. We infer that it coincides with $\hat{\rho}_T\alpha$, thereby completing the proof. □

3. Proof of Theorem A and Corollary B

Lemma 3.1. *For distinct edges $b_i, b_j \in \Gamma_1$, let $\mathcal{T}_{ij}$ be the set of ρ-spanning trees such that $\langle \bar{T}(b_i), b_j\rangle \neq 0$. Then*

$$\sum_{T\in\mathcal{T}_{ij}} w_T\langle \bar{T}(b_i), b_j\rangle_R = \sum_{U\in\mathcal{T}_{ji}} w_U\langle b_i, \bar{U}(b_j)\rangle_R\,.$$

Proof. From the definition of the weights, have

$$\frac{r_j w_T}{\hat{\rho}_T} = \frac{r_i w_U}{\hat{\rho}_U}\,. \tag{8}$$

Recall that $\langle \bar{T}(b_i), b_j\rangle_R = r_j\langle \bar{T}(b_i), b_j\rangle$. Using Eq. (8) and Proposition 2.7, we infer

$$w_T\langle \bar{T}(b_i), b_j\rangle_R = w_U\langle b_i, \bar{U}(b_j)\rangle_R\,.$$

Now sum up over all $T \in \mathcal{T}_{ij}$. □

Proof of Theorem A. Consider the operator $F := \sum_T w_T \bar{T}$, where the sum is over all ρ-spanning trees of Γ. For any pair of edges b_i and b_j of Γ we have

$$\begin{aligned}
\langle \sum_T w_T \bar{T}(b_i), b_j \rangle_R &= \sum_{T \in \mathcal{T}_{ij}} w_T \langle \bar{T}(b_i), b_j \rangle_R \\
&= \sum_{U \in \mathcal{T}_{ji}} w_U \langle b_i, \bar{U}(b_j) \rangle_R \qquad \text{by Lemma 3.1 ,} \\
&= \langle b_i, \sum_U w_U \bar{U}(b_j) \rangle_R \\
&= \langle b_i, \sum_T w_T \bar{T}(b_j) \rangle_R
\end{aligned}$$

Hence F is self-adjoint in the modified inner product.

If $z \in Z_1(\Gamma; \rho)$, then using Corollary 2.5, we have

$$F(z) = (\sum_T w_T)\bar{T}(z) = (\sum_T w_T) z =: \Delta z$$

Consequently, $(1/\Delta)F$ restricts to the identity on $Z_d(X; \rho)$. As $(1/\Delta)F$ is self-adjoint, it is the Hermitian projection in the modified inner product. □

Proof of Corollary B. Let z be the Hermitian projection of $R^{-1}\mathbf{V}$ in the modified inner product. Then $R^{-1}\mathbf{V} - z \in B^d_R(\Gamma; \rho)$, i.e.,

$$0 = \langle R^{-1}\mathbf{V} - z, z' \rangle_R = \langle \mathbf{V} - Rz, z' \rangle$$

for all $z' \in Z_d(X; \rho)$. Hence, $\mathbf{V} - Rz \in B^d(\Gamma; \rho)$. The uniqueness of z is a consequence of the fact that $B^d(\Gamma; \rho)$ is the orthogonal complement to $Z_d(X; \rho)$ in the standard inner product.

The proof of the last part is given by direct calculation using the self-adjointness of the operator $\sum_T w_T \bar{T}$:

$$\begin{aligned}
\langle z, b \rangle &= \tfrac{1}{r_b} \langle z, b \rangle_R \,, \\
&= \tfrac{1}{r_b} \langle \tfrac{1}{\Delta} \textstyle\sum_T w_T R^{-1}\mathbf{V}, b \rangle_R \,, \\
&= \tfrac{1}{\Delta} \sum_T \tfrac{w_T}{r_b} \langle R^{-1}\mathbf{V}, \bar{T}(b) \rangle_R \,, \\
&= \tfrac{1}{\Delta} \sum_T \tfrac{w_T}{r_b} \langle \mathbf{V}, \bar{T}(b) \rangle \,.
\end{aligned}$$

□

4. Proof of Theorem C

The proof of Theorem C is essentially the same as the proof of [CCK, th. C]. We will outline the essential steps. The first step is to show that

$$\det(\partial\partial_R^*\colon C_0(\Gamma;\rho)\to C_0(\Gamma;\rho)) = \gamma\sum_T w_T \tag{9}$$

where T ranges over all ρ-spanning trees, and the pre-factor γ is to be determined. This step follows, *mutatis mudandis*, by the proof of [CCK, prop. 4.2].

The second and final step is to compute the pre-factor γ and to show that it equals 1. Let use write $W\colon \Gamma\to\mathbb{R}$ for the function given by $b\mapsto \ln r_b$. Then we have $R = e^W$. Fix a ρ-spanning tree T. Then as in [CCK, defn. 5.1], we say that W is *good* if

$$W_b > \sum_{b'\in T_1} W_\alpha - k\min_{b'\in T_1} W_{b'} \qquad \text{for any } b\in\Gamma_1\setminus T_1,$$

where k is the number of edges of Γ.

Set $\mathcal{L}_R = \partial\partial_R^*$ and similarly define $\mathcal{L}_R^T\colon C_0(T;\rho)\to C_0(T;\rho)$ by replacing Γ with T. Let $\beta\in R_+$ represent inverse temperature. Then by the argument which proves [CCK, prop. 5.2], we have

$$\lim_{\beta\to\infty}\frac{\mathcal{L}_{R_\beta}^T}{\mathcal{L}_{R_\beta}} = 1 \tag{10}$$

where $R_\beta := e^{\beta W}$.

Substituting βW for W in Eqn. (9), taking the low temperature limit $\beta\to\infty$, substituting the relation (10) and some minor rewriting, we deduce

$$\det(\mathcal{L}_R^T) = \gamma w_T.$$

Note that $\mathcal{L}_R^T = \partial_T e^{-W}\partial_T^*$, and by definition of w_T, we have $\det e^{-W} = \hat\rho_T^{-1} w_T$. Consequently,

$$\det(\mathcal{L}_R^T) = \hat\rho_T^{-1} w_T \det(\partial_T\partial_T^*).$$

It follows that

$$\gamma = \hat\rho_T^{-1}\det(\partial_T\partial_T^*).$$

Theorem C is then a consequence of the following.

Lemma 4.1. *For any ρ-spanning tree T, we have*

$$\det(\partial_T\partial_T^*) = \hat\rho_T\,.$$

Hence, $\gamma = 1$.

Proof. Clearly both sides of the equation factor as a product of over the connected components of T. So if T^α is a component of T, it will suffice to show

$$\det(\partial_{T^\alpha}\partial^*_{T^\alpha}) = \hat{\rho}_{T^\alpha}\,.$$

This last statement can be proved in a number of ways. For example, Kenyon [Ke] proves it using an interpretation of the determinant as a summation of cycles over the symmetric group. We will give a proof using gauge invariance.

The *gauge group* G of Γ is the group of functions $\Gamma_0 \to U(1)$ with respect to pointwise multiplication. It is convenient in what follows to set $g_v = g(v)$ for a vertex v. Then G acts on line bundles according to the rule

$$g \cdot \rho(b) = g_{d_0(b)} g^*_{d_1(b)} \rho_b\,.$$

Set $\rho^g = g \cdot \rho$. To distinguish between boundary operators, we write ∂ for the boundary operator associated with ρ, and ∂_g for the one associated with ρ^g. Define an action

$$G \times C_0(\Gamma;\rho) \to C_0(\Gamma;\rho)$$

by $g \cdot v = g_v v$, for $v \in \Gamma_0$.

It is then straightforward to check that for $g \in G$ we have

$$\partial_g \partial^*_g = g \partial \partial^* g^{-1}.$$

In particular, $\det(\partial_g\partial^*_g) = \det(\partial\partial^*)$.

Write $T^\alpha = A \cup b$, where A is tree in the classical sense. We claim that there is a gauge $g \in G$ such that $\partial_g(e) = 1$ for $e \in A_1$. To find g we need to know that the system of equations

$$g_{d_0(e)} g^*_{d_1(e)} \rho_e = 1, \qquad e \in A \tag{11}$$

admits a solution. If we fix a vertex $i \in A_0$, we can set $g_i = 1$. Then for any edge ij of A which connects i to j, we set $g_j = \rho^s_{ij}$ with $s = \pm 1$ according as to whether ij points inward towards i or not. Consider a vertex $k \neq i$ such that jk is an edge of A. We set $g_k = \rho^s_{jk} g_j$, where in this instance s is ± 1 according as to whether the edge jk points towards j or not. Continuing in this fashion, we obtain a solution to the system (11). For any vertex i not in A we set $g_i = 1$. With respect to our choice of g, inspection shows that ρ^g_b is the holonomy with respect to ρ around the unique circuit of $A \cup b$ which is oriented in the direction of b. Hence, we can without loss in generality assume that the original line bundle ρ is trivial on every edge other than b, and we are reduced to proving that $\det(\partial\partial^*) = (\rho_b - 1)(\rho^*_b - 1)$, where ∂ is the boundary operator for $T^\alpha = A \cup b$.

The columns of the matrix associated with ∂ are given by the edges of $A \cup b$ and the rows are given by the vertices. An edge e of $A \cup b$ is said to be *loose* if it is attached to a vertex i such that no other edge of $A \cup b$ is attached to i. If e is loose, then the e-th column of ∂ has exactly two non-zero entries which are ± 1 and these are of opposite sign. We infer that the determinant of ∂ remains unchanged when we remove the edge e and the vertex i from $A \cup b$. Iterating this procedure, we may assume without loss in generality that $A \cup b$ has no free edges. This means $A \cup b$ is a circuit. The determinant of ∂ in this case is easy to compute and is given by $\pm(\rho_b - 1)$. Hence the determinant of $\partial^* \partial$ is $(\rho_b - 1)(\rho_b^* - 1)$. □

References

[B] Bollobás, B.: Modern graph theory. Graduate Texts in Mathematics, 184. Springer-Verlag, New York, 1998.

[CCK] Catanzaro, M.J., Chernyak, V.Y., Klein, J.R.: Kirchhoff's theorems in higher dimensions and Reidemeister torsion. *arXiv preprint 1206.6783*

[E] Eckmann, B.: Harmonische Funktionen und Randwertaufgaben in einem Komplex. *Comment. Math. Helv.* **17** (1945), 240–255.

[F] Forman, R.: Determinants of Laplacians on graphs. *Topology* **32** (1993), 35-46

[Ke] Kenyon, R.: Spanning forests and the vector bundle Laplacian *Ann. Probab.* **39** (2011), 1983-2017

[NS] Nerode, A., Shank H.: An algebraic proof of Kirchhoff's network theorem, *Amer. Math. Monthly* **68**, (1961) 244–247.

[R] Roth, J.P.: An application of algebraic topology to numerical analysis: on the existence of a solution to the network problem. *Proc. Nat. Acad. Sci. U.S.A.* **41** (1955), 518–521

Department of Mathematics, Wayne State University, Detroit, MI 48202

E-mail address: mike@math.wayne.edu

Department of Chemistry, Wayne State University, Detroit, MI 48202

E-mail address: chernyak@chem.wayne.edu

Department of Mathematics, Wayne State University, Detroit, MI 48202

E-mail address: klein@math.wayne.edu